



Know! Alcohol's Female Effect

April is Alcohol Awareness Month. In this Know! Parent Tip we focus on alcohol's effects specific to young females.

Many adults assume that underage drinking mostly involves boys. That assumption however, is false. In fact, girls have not only caught up to boys when it comes to drinking, but in many cases have surpassed them. According to results from the 2012 National Survey on Drug Use and Health, among youth aged 12 to 17, the percentage of females who were current drinkers (13.2 percent) was higher than their male counterparts (12.6 percent).

Research also shows that while alcohol use among boys and girls is just about equal, alcohol's impact on their bodies and brains are not.

Females are more vulnerable to alcohol's effects than males, and here's why:

- Females have less water in their bodies to help dilute the alcohol in the bloodstream;
- Females absorb alcohol at a slower rate;
- Females naturally produce less *alcohol dehydrogenase* (a gastric enzyme that breaks down ethanol in the stomach – that otherwise is toxic).

What this means is that a female and male of the same size and weight can drink the same amount of alcohol and yet the female will have a higher concentration of alcohol in her blood. It also means that females who go "drink-for-drink" with males are likely to become intoxicated more quickly and are more susceptible to alcohol poisoning.

Alcohol can be damaging to the developing adolescent brain, regardless of gender. However, females are more sensitive to alcohol-induced brain damage than males. Research has found that females who drink heavily for long periods of time may experience a reduction in their *corpus callosum* (a band of nerves deep within the brain that connect the left and right hemispheres of the brain to communicate and coordinate one's activities), whereas males do not.

As females age, the differences remain. As adults, women struggle more than men to break down fatty acids in the body, therefore resulting in a higher number of women who develop liver diseases after comparatively shorter periods of heavy drinking than men. But even younger women in their late teens and early 20s who chronically abuse alcohol are at increased risk for ulcers and other gastrointestinal problems.

Talking regularly with kids about the dangers of alcohol, tobacco and other drugs reduces their risk of using.

Know! urges you to encourage other parents to join Know!.

[Click here for the Know! Parent Tip Sign-Up Page.](#)

Know! is a program of:



*Lifetime Prevention
Lifetime Wellness*

Drug Free
Action Alliance
6155 Huntley Rd.
Suite H
Columbus, Ohio
43229



Link to Facebook
Page

Underage drinking is damaging and dangerous. As an important influencer in your students' lives, teachers are encouraged to talk early and often, sharing clear anti-use messages with your female and male students alike. It is important to include in your talks, especially with the girls, the fact that drinking impacts females more intensely, so that they are better armed to make informed, healthy decisions surrounding alcohol, as teens and into their adult years.

Sources: [2012 National Survey on Drug Use and Health: Summary of National Findings. Girls, Women and Alcohol: Making Informed Choices resource from Healthy Child Manitoba.](#) [NYU Langone Medical Center – The Child Study Center: Girls and Alcohol.](#)